

Paracoccidioidomycosis disease (Lutz-Splendore-Almeida) – clinical manifestations

Paracoccidioidomicose (doença de Lutz-Splendore-Almeida) – manifestações clínicas

Alexandre Vasconcellos Alvim Ambrósio¹, Camila Cristiane Silva Camelo¹, Carolina Venâncio Barbosa¹, Fernanda Gomes Tomazatti¹, Flávia Araújo de Souza Brazões¹, Juliana Márcia Veloso¹, Gregório Victor Rodrigues¹, Lucas Fonseca Rodrigues¹, Pedro Igor Daldegan de Oliveira¹, Raíza Almeida Aguiar¹, Valdirene Silva Siqueira¹, Victor Bastos Jardim¹, Victoria Almeida Correa Gontijo¹, Samuel Gonçalves da Cruz², Weverton César Siqueira², Enio Roberto Pietra Pedrosa³

DOI: 10.5935/2238-3182.20140019

ABSTRACT

Paracoccidioidomycosis has polymorphic clinical features with lesions located in the skin and mucous membranes, as well as involvement of various organs and systems, as is potentially capable of causing death and serious sequelae. It should be included in the differential diagnosis of granulomatous diseases in endemic areas, including Brazil, so that it is recognized early, for more convenient treatment as to prevent progression with sequelae or premature death.

Key words: Paracoccidioidomycosis; Mycosis; Paracoccidioides.

RESUMO

A paracoccidioidomicose possui clínica polimórfica, com manifestações localizadas em pele e mucosas até comprometimento de vários órgãos e sistemas, potencialmente capaz de provocar sequelas graves e morte. Deve ser incluída no diagnóstico diferencial das doenças granulomatosas, em áreas endêmicas, como ocorre no Brasil, para que seja reconhecida com precocidade, tratada convenientemente e evitada sua evolução para sequelas e morte prematura.

Palavras-chave: Paracoccidioidomicose; Micose; Paracoccidioides.

INTRODUCTION

The importance of paracoccidioidomycosis (PCM) relates to social and economic costs not only derived from the active disease in individuals in the most productive phase of their lives, but from sequelae, which represent reasons for work incapacitation. The usual evolution of paracoccidioidomycosis without specific therapeutic intervention is to the death.^{1,2}

This disease constitutes one of the neglected diseases by all funding agencies for its study. Despite of being endemic in Brazil, it is not always considered in the differential diagnosis of diseases that are localized with lymphadenomegaly, infiltration, and mucocutaneous vegetative-nodules, pulmonary fibrosis; or systemic with different involvement, from adrenal insufficiency to severe neurological.^{3,4}

Whenever the differential diagnosis includes tuberculosis and lymphoma, the possibility of PCM must be considered, which means valuing the disease's clinical epidemiology (geography of diseases).⁵⁻¹⁷

Submitted: 1/16/2014
Approved: 2/10/2014

Institution:
Medical School at UFMG
Belo Horizonte, MG – Brazil

Corresponding Author:
Enio Roberto Pietra Pandey
E-mail: enio@medicina.ufmg.br

CLINICAL MANIFESTATIONS

PCM can manifest itself in different forms: acute-sub-acute, regressive, evolutionary, or chronic and slowly progressive. Several organs and systems can be affected (Table 1). The characterization of its various forms is made as a function of the patient's age; illness form of presentation and duration; clinical manifestations; associated diseases and aggravating factors, general and nutritional state, thorax telerradiography, response to the cutaneous test with paracoccidioidin, and serum levels of antibodies anti *P. brasiliensis* (double immunodiffusion reaction in agar gel).^{11, 14,16-27}

Table 1 - Classification of the clinical forms of paracoccidioidomycosis

1. Paracoccidioidomycosis infection
2. Paracoccidioidomycosis disease
2.1. Regressive form
2.2. Progressive form
2.2.1. Acute or sub-acute (youth form), with:
a) Superficial adenomegaly (moderate and severe)
b) Superficial adenomegaly (moderate and severe)
c) Bone involvement (severe)
d) Other clinical manifestations (moderate or severe)
2.2.2. Chronic (adult form): light, moderate, severe
2.2.3. With sequelae

Therefore, the host-fungus interaction relates to the type and intensity of the immune response, virulence, and fungus inhalation, which determines how the initial infection may present varying courses, either for the acute-sub-acute or chronic and slowly progressive, or, more commonly, be resolved but leaving a latent focus that can reactivate (endogenous reactivation) and subsequently generate the chronic form (Table 2).¹⁶⁻²⁵

Table 2 - Classification according to the seriousness of the acute, sub-acute, and chronic forms of paracoccidioidomycosis

Forms		Clinical alterations						
		Lymphnodes	H-E	General state	Lungs and/or skin	Other organs	PCD	Ac
Acute-sub-acute	M	Inflammatory, non-suppurative	P/A	P/A	A	A	35 mm	B/M
	G	Tumoral or suppurative	P	P	P	P	5 mm	E
Chronic	M	Inflammatory, non-suppurative	P/A	P/A	A/P	A	≥10 mm	B
	G	Tumoral or suppurative	P	P	P	P	<5 mm	E

M: moderate; G: severe; P: present; A: absent; E: high; B: low; M: moderate; L: lymphadenomegaly; H-E: hepatosplenomegaly; PCD: reaction to paracoccidioidin; Ac: serum levels of antibodies by immunodiffusion.

Regressive form

The host initial invasion by Paracoccidioidis is generally asymptomatic with possible varied clinical manifestations, from discrete to nonspecific, such as from asthenia, anorexia, evening febricula, to febrile syndrome, cough, hemoptoic, and chest pain. It resembles tuberculosis in its phase of primary complex, and constitutes a most benign form of PCM.

Its total regression can coexist with the presence of the fungus, as the quiescent form, in lymph node or lung. It can leave the skin reaction to paracoccidioidin in the memory.^{11,13-16}

Acute -Sub-acute form

From the location of the initial invasion or quiescent focus, the Paracoccidioidis may spread via bronchogenic, lymphatic, or hematogenous, and in 34% of cases, it is manifested as a disease of systemic acute sub-acute evolution with involvement of mononuclear-phagocyte system.²⁸⁻³⁴

It preferentially affects children, adolescents, and young adults up to 30 years old, becoming the youth age form and considered of moderate to intense gravity.²⁸⁻³⁴

The evolution of PCM is usually of short duration, rapidly progressive, debilitating, with the development of: asthenia, anorexia, intense weight loss capable of causing cachexia; diffuse lymphadenomegaly, with necrosis and suppuration, expressing itself as cutaneous or intra-abdominal abscesses, cutaneous fistulas with drainage of purulent material, and cutaneous areas of extensive destruction (strophulus); osteomyelitis; intestinal ulcerations; hepatosplenomegaly, and hypo or aplasia of bone marrow.

The intra-abdominal lymphadenomegaly can form bulky masses that compress different structures and determine various clinical syndromes such as biliary obstruction (cholestasis), pancreatic duct (pancreatitis), thoracic duct (chylous ascites), ureters (pyelonephritis, acute renal failure), and intestines (malabsorption syndrome, acute abdomen). Mucosal (20%) and lung (10%) alterations are infrequent. There may be increased body temperature, which represents a sign of seriousness. Pulmonary involvement, contrary to what occurs in the chronic form, is rare.

The acute or sub-acute clinical manifestations can also occur in adults, along with lymphadenomegalies that disinfect, intense eosinophilia, and serum precipitating antibodies (double immunodiffusion reaction) at higher levels than those observed in the chronic form.⁵⁻³⁴

Chronic form

It generally affects people after the fourth decade of life, slowly and progressively, from the site of Paracoccidioidis invasion or its quiescent focus (latent), which can reactivate later (endogenous reactivation), lasting more than six months.^{11, 14, 16, 35-81}

Its clinical manifestations are variable, from light to even negligible to moderate and severe, with progressive impairment of the general condition with significant weight loss. The most common clinical alterations arise from the involvement of the skin, mucous membranes of the upper respiratory tract and mouth, lymph nodes, and lungs.

The main clinical manifestations are: anterior and posterior cervical lymphadenomegalies (79%), submandibular, submentovertebral, supraclavicular, axillary, inguinal, and intra-abdominal (expressed through acute abdomen, tumor masses, jaundice by biliary extra-hepatic compression, chylous ascites, or intestinal malabsorption); weight loss (72%); asthenia/hypodinamia (65%); mucocutaneous paleness (62%); fever (51%); cough (50%); dysphonia, odynophagia, dysphagia, oropharyngeal ardor, nasal obstruction, epistaxis; vegetative ulcers in the mouth, throat, and nose.

The cutaneous manifestations present varied forms, with single or multiple lesions, infiltrative, vegetative verruciform, expressed primarily as papules, plaques, nodules, vegetative, or ulcers. The presence of vegetative lesions is frequent, ulcerated and painless in the oral and palate cavities (Figure 1), tongue (Figure 2), gum and lip (Figures 3 and 4), and nasal,

in the pharynx, and larynx, including with satellite adenopathy (Figures 4, 5, and 6) (cervical, axillary, and inguinal). The consistency of lesions is, usually, hardened due to fibrosis in the chronic granulomatous process. The main locations are the face, upper and lower limbs, and trunk. It can evolve with secondary bacterial infections. Purely infiltrative lesions are rare with sarcoid pattern, usually with few fungi observed in microscopy, which makes the diagnosis difficult.⁸⁻²³



Figure 1 - Moriforme lesion in the patient's palate with multifocal chronic form. Patient assisted at the PCM Reference Center in the General Hospital from UFMG.




Figure 2 - Ulcer and painful lesion in tongue with sialorrhea. Patient assisted at the PCM Reference Center in the General Hospital from UFMG.

Lymphadenomegaly (Figures 5 and 6) can present inflammatory non-suppurative features (less than 2 cm in diameter), tumoral (more than 2 cm in diameter), and suppurative (floating or fistulization).^{11-14, 16-23}



Figure 3 - Two oral lesions, one in the lower lip and one in the gum, with tooth loss; ulcerated, hardened, granulomatous and elevated edges. Patient assisted at the PCM Reference Center in the General Hospital from UFMG.



Figure 4 - Skin lesions in erythematous crusted papules and nodules on the face, in the multifocal chronic form, before (A) and after (B) treatment. Patient assisted at the PCM Reference Center in the General Hospital from UFMG.



Figure 5 - Cervical lymphadenomegalies and hardened papules in the malar region in the multifocal chronic form. Patient assisted at the PCM Reference Center in the General Hospital from UFMG.



Figure 6 - Inguinal bilateral lymphadenomegaly in patient with the PCM multifocal chronic form. Patient assisted at the PCM Reference Center in the General Hospital from UFMG.



Figure 7 - Skin lesions in the multifocal chronic form. Hardened nodules with elevated edges in the trunk (A) and legs (B). Erythematous papules and plaques, crusted and diffused on the face (C). Ulcerated lesion on the glands (D). Patient assisted at the PCM Reference Center in the General Hospital from UFMG.

The oropharynx (Figure 1) can present hyperemia, edema, and a moriforme aspect, infiltration or vegetating, or even ulcerated, teeth can be softened, and drilling of the hard palate can be observed.

Respiratory alterations are also of varying intensity, from mild to severe, with the physical examination of the thorax negligible in most patients, characterized by clinical-radiological semiological dissociation in which light symptoms contrasts with intense radiological alterations in the pleuropulmonary fields.

Adrenals, digestive systems, bones, and central nervous systems (CNS) alterations can still be observed.

The involvement of the adrenal occurs in 50% of cases and can determine medullar insufficiency, similar to tuberculosis lesions, as Addison's syndrome, evolving from an insidious form with asthenia, anorexia, weight loss, postural hypotension, fainting, dizziness, hyperpigmentation of the skin and mucous membranes, nausea and vomiting, reduced sexual potency and libido, and blood eosinophilia. The adrenal gland does not always retrieve its function after the specific treatment for PCM.

The CNS can be affected in its parenchyma or in the meninges in almost 25% of the cases. It can evolve as a pseudotumor or meningeal form (diffuse or localized lesions, with more frequent involvement of the base of the skull). Its evolution is dragged and resembles the tuberculous meningitis, with which it makes for a differential diagnosis.⁷⁷⁻⁸¹ The parenchymal form is associated with: progressive intracranial hypertension installation, signs of focal dysfunction, motor or sensory, of language, and ataxia alterations. It can be associated to focal or generalized convulsions and papilla edema. Spinal cord alterations may lead to paresthesia, anesthesia, and lower limb weakness, urinary and fecal incontinence and neurogenic bladder, with episodes of urinary retention.

Digestive alterations are varied and non-specific in most cases, presenting itself as a tumoral mass, with malabsorption and chronic diarrhea.

Bone alterations (20%) can be asymptomatic and revealed by lytic radiological lesions without perifocal reaction, with possible periosteal involvement, light and with sharp edges. Bone marrow is engaged mainly in the form of acute or sub-acute and rarely in the chronic forms, manifesting itself in the peripheral blood in combination with anemia, leukopenia, and thrombocytopenia.^{11, 14, 16, 28-34, 39-76}

The least common manifestations are genitourinary (Figure 7 D), characterized by the involvement

of the epididymis, testis, and prostate, in which complaints of dysuria, alguria, or hematuria dominate; in the thyroid; and eyes, which can be unilaterally altered, with lesions on the eyelids (papule, vegetation, ulcer), conjunctiva, anterior uvea, or choroid.³⁵⁻⁸¹

The average time until death in patients who evolved spontaneously with PCM is from five to eight months, comparable to visceral leishmaniasis and aggressive malign neoplasias.^{1, 2, 6, 7, 11-16}

The forms with sequelae are frequent; the lung forms stand out with the predominance of fibrosis and emphysema; Addison's syndrome; neurological lesions; and disfiguring skin forms (Table 2).³⁵⁻⁸¹

REFERENCES

1. Moreira APV. Paracoccidioidomycose: histórico, etiologia, epidemiologia, patogênese, formas clínicas, diagnóstico laboratorial e antígenos. *Bol Epidemiol Paulista*. 2008 mar; 5(51).
2. Marques SA. Paracoccidioidomycose: atualização epidemiológica, clínica e terapêutica. *An Bras Dermatol*. 2003; 78(2):135-50.
3. Souza W. (coordenador). Doenças negligenciadas. Rio de Janeiro: Academia Brasileira de Ciências; 2010.
4. Martinez R. Paracoccidioidomycosis: The dimension of the problem of a neglected disease. *Rev Soc Bras Med Trop*. 2010; 43:480.
5. Fava Netto C, Castro RM, Gonçalves AP, Dillon NL. Ocorrência familiar de blastomicose sul-americana: a propósito de 14 casos. *Rev Inst Med Trop São Paulo*. 1965; 7(6):332-6.
6. Pereira RM, Tresoldi AT, Silva MTN, Bucaretych F. Fatal disseminated paracoccidioidomycosis in a two-year-old child. *Rev Inst Med Trop S Paulo*. 2004; 46(1):37-9.
7. Araújo SA, Prado LG, Veloso JM, Pedroso ERP. Case of recurrent Paracoccidioidomycosis: 25 years after initial treatment. *Braz J Infect Dis*. 2009; 13(5): 394.
8. Motta LC. Granulomatose paracoccidioidica (blastomicose brasileira). *Ann Fac Med S Paulo*. 1936; 12:407-26.
9. Bopp C. Algumas considerações sobre a micose de Lutz no Rio Grande do Sul. *An Fac Med P Alegre*. 1955; 26:97-106.
10. Lacaz CS. Paracoccidioidomycose. In: Lacaz CS, Porto E, Martins JEC. *Micologia Médica*. 8. ed., São Paulo: Sarvier; 1991. p. 248-97.
11. Bummer E, Castaneda E, Restrepo A. Paracoccidioidomycosis: an update. *Clin Microbiol Rev*. 1993; 6:89.
12. Machado Filho J, Miranda JL. Considerações relativas à blastomicose sul-americana: localizações, sintomas iniciais, vias de penetração e disseminação em 313 casos consecutivos. *O Hospital*. 1960; 58:99-137.
13. Machado Filho J, Miranda JL. Considerações relativas à blastomicose sul-americana. *O Hospital*. 1961; 60(4):21-62.
14. Mendes RP. Paracoccidioidomycose. In: Rocha MOC, Pedroso ERP. *Fundamentos em Infectologia*. Rio de Janeiro: Rubio; 2009. p. 945-94.

15. Montenegro MR. Formas clínicas da paracoccidioidomicose. *Rev Inst Med Trop S Paulo*. 1986; 28:203-4.
16. Restrepo A, Tobon AM, Agudelo CA. Paracoccidioidomycosis. In: Hospenthal DR, Rinaldi MG. (editors). *Diagnosis and treatment of human mycoses*. Totowa, NJ: Humana Press; 2008. p. 331.
17. Rizzon CFC, Severo LC, Porto NC. Paracoccidioidomicose, estudo de 82 casos observados em Porto Alegre, RS. *Rev AMRIGS*. 1980; 24(1):15-7.
18. Padilha GA. Paracoccidioidomicose: quadro clínico como expressão da imunopatologia. *Anais Brasileiros Dermatol*. 1996; 71(5):1996.
19. Paniago AMM, Aguiar JIA, Aguiar ES, Cunha RV, Pereira GROL, Londero AT, *et al*. Paracoccidioidomicose: estudo clínico e epidemiológico de 422 casos observados no estado de Mato Grosso do Sul. *Rev Soc Bras Med Trop*. 2003; 36(4):455-9.
20. Pedrosa ERP, Veloso JMR, Prado LGR. Saúde pública e epidemiologia: a paracoccidioidomicose em pacientes atendidos no Hospital das Clínicas (HC-UFMG). In: III Congresso Mineiro de Infectologia, 2008, Belo Horizonte. A paracoccidioidomicose em pacientes atendidos no Hospital das Clínicas (HC-UFMG). Belo Horizonte: HC-UFMG; 2008.
21. Gontijo CCV, Prado RS, Neiva CLS, Freitas RM, Prado FLS, Pereira ARA, *et al*. A paracoccidioidomicose em pacientes atendidos no Hospital das Clínicas da UFMG (HC-UFMG). *Rev Med Minas Gerais*. 2002; 13(4):231-3.
22. Lima FXP. Contribuição ao estudo clínico e terapêutico da blastomicose sul-americana. [Dissertação]. São Paulo: Faculdade de Medicina da Universidade de São Paulo; 1952.
23. Londero AT, Del Negro G. Paracoccidioidomicose. *J Pneumol*. 1986; 12:41-60.
24. Louzada A. Blastomicose e tuberculose. *Med Cir*. 1954; 16:38-42.
25. Morejón KM, Machado AA, Martínez R. Paracoccidioidomycosis in patients infected with and not infected with human immunodeficiency virus: a case-control study. *Am J Trop Med Hyg*; 2009; 80:359.
26. Franco M, Montenegro MR, Mendes RP, Marques SA, Dillon NL, Mota NG. Paracoccidioidomycosis: a recently proposed classification of its clinical forms. *Rev Soc Bras Med Trop*. 1987; 20(2):129-32.
27. Shikanai Yasuda MA, Telles Filho FO, Mendes RP. Guidelines in paracoccidioidomycosis. *Rev Soc Bras Med Trop*. 2006; 39:297.
28. Barba MF, Marques HHS, Scatigno Neto A, Aquino MZ, Vitule LF, Barbato AJG, *et al*. Paracoccidioidomicose na infância: diagnóstico por imagens – relato de caso. *Radiol Bras*. 1993; 26(2):87-90.
29. Pereira RM, Bucarechi F, Barison Ede M, Hessel G, Tresoldi AT. Paracoccidioidomycosis in children: clinical presentation, follow-up and outcome. *Rev Inst Med Trop São Paulo*. 2004; 46:127-31.
30. Pereira RM, Bucarechi F, Barison EM, Hessel G, Tresoldi AT. Paracoccidioidomycosis in children: clinical presentation, follow-up and outcome. *Rev Inst Med Trop São Paulo*. 2004; 46(3):127.
31. Castro RM, Del Negro G. Particularidades clínicas da paracoccidioidomicose na criança. *Rev Hosp Clin*. 1976; 31(3):194-8.
32. Cruz MLS, Souza MJ, Glaura MFT, Machado ES, Rios-Gonçalves AJ. Paracoccidioidomicose: a propósito de mais um caso da forma infantil. *Arq Bras Med*. 1991; 65(2):102-4.
33. Ferreira MS. Contribuição para o estudo clínico-laboratorial e terapêutico da forma juvenil da paracoccidioidomicose. *Rev Pat Trop*. 1993; 22(2):267-406.
34. Nogueira MG, Andrade GM, Tonelli E. Clinical evolution of paracoccidioidomycosis in 38 children and teenagers. *Mycopathol*. 2006; 161:73.
35. Baldó JI. Paracoccidioidomycosis pulmonar. *Rev Sanid Assist Soc*. 1953; 18:163-76.
36. Machado Filho J, Carvalho MCM. Considerações em torno das localizações pulmonares da paracoccidioidomicose brasileira. *Rev Bras Tuber*. 1952; 9/10:9-41.
37. Freitas RM, Prado R, Prado FL, Paula IB, Figueiredo MT, Ferreira CS, *et al*. Pulmonary paracoccidioidomycosis: radiology and clinical-epidemiological evaluation. *Rev Soc Bras Med Trop*. 2010; 43(6):651-6.
38. Furtado T. Comprometimento pulmonar na blastomicose sul-americana. *Rev Assoc Méd Minas Gerais*. 1952; 3:49-57.
39. Gonçalves AJR, Guerra S, Jesus PLL. Paracoccidioidomicose aguda escavada associada a carcinoma da paratireóide. *F Med (BR)*. 1980; 81(6):623-6.
40. Costa MAB, Carvalho TN. Manifestações extrapulmonares da paracoccidioidomicose. *Radiol Bras*. 2005; 38(1):45-55.
41. Fialho F. Localizações viscerais da micose de Lutz. *Bol Acad Nac Med*. 1962; 134:18-23.
42. Moraes CR. Calcificações intra-abdominais na blastomicose sul-americana. *Rev Inst Med Trop S Paulo*. 1971; 13(6):428-32.
43. Andrade ALSS. Paracoccidioidomicose linfático-abdominal: contribuição ao seu estudo. *Rev Pat Trop*. 1983; 12(2):165-256.
44. Rocha A. Blastomicose sul-americana: caso de forma linfático-abdominal pseudotumoral. *O Hospital*, 69(4):157-163, 1966.
45. Gonçalves AJR, Pinto AMM, Mansur Filho J, Silva Filho MC. Paracoccidioidomicose – formas linfáticoabdominal pseudotumoral e disseminada com abdômen agudo e manifestações neurológicas. *Arq Bras Med*. 1983; 51(1):7-10.
46. Andrade DR. Hipoproteinemia em pacientes com paracoccidioidomicose do tubo digestivo e sistema linfático abdominal: revisão de casos de necropsia e apresentação de um caso com perda protéica digestiva. *Rev Hosp Clin*. 1976; 31(3):174-9.
47. Barbosa W, Daher R, Oliveira AR. Forma linfático-abdominal da blastomicose sul-americana. *Rev Inst Med Trop S Paulo*. 1968; 10(1):16-27.
48. Silva AL, Giacomini RT, Silva BD. Paracoccidioidomicose ganglionar calcificada. *Rev Soc Bras Med Trop*. 1991; 24(4):253-5.
49. Angulo-Ortega A. Calcifications in paracoccidioidomycosis: are they the morphological manifestation of subclinical infections? In: Pan American Symposium on Paracoccidioidomycosis, 1 ed. Medellin, 1971. PAHO Scient Publ. 1972; 254:129-33.
50. Rocha A. Blastomicose sul-americana ganglionar primitiva com calcificações abdominais. *O Hospital*. 1966; 70(6):195-206.
51. Rocha A. Abdome agudo por colecistite blastomictica: relato de um caso. *Rev Goiana Med*. 1980; 26:63-9.
52. Gabellini GC, Martinez R, Ejima FH, Saldanha JC, Módena JLP, Velludo MAL, *et al*. Paracoccidioidomicose do estômago: relato de caso e considerações sobre a patogenia da lesão. *Arq Gastroenterol*. 1992; 29(4):147-52.

53. Gonçalves AJR. Aneurisma blastomicótico da aorta abdominal. Relato de caso e revisão de literatura. *Rev Med Hosp Serv Est.* 1977; 29(3):119-24.
54. Gonçalves AJR, Terra GMF, Rozembaum R, Londero AT. Paracoccidioidomicose com enterocolite em criança de 12 anos. *J Bras Med.* 1996; 70(4):50-4.
55. Martinez R, Rossi MA. Paracoccidioidomicose duodenal com sangramento digestivo. *Rev Inst Med Trop S Paulo.* 1984; 26:160-4.
56. Martinez R, Meneghelli UG, Dantas RO, Fiorillo AM. O comprometimento gastrointestinal na blastomicose sul-americana (paracoccidioidomicose). Estudo clínico, radiológico e histopatológico. *Rev Ass Med Brasil.* 1979; 25(1):31-4.
57. Martinez R. Manifestações digestivas da paracoccidioidomicose. In: Del Negro G, Lacaz CDAS, Fiorillo AM. *Paracoccidioidomicose: blastomicose sul-americana.* São Paulo: Sarvier; 1982. p. 483-91.
58. Fiorillo AM, Martinez R, Moraes CR. Lesões do aparelho digestivo. In: Del Negro G, Lacaz CS, Fiorillo AM. *Paracoccidioidomicose: blastomicose sul-americana.* São Paulo: Sarvier; 1982. p. 189-93.
59. Barbosa W, Andrade ALSS. Hepatic involvement in paracoccidioidomycosis. *Rev Pat Trop.* 1983; 12(3):295-305.
60. Bocallandro I, Albuquerque FJM. Icterícia e comprometimento hepático na blastomicose sul-americana. A propósito da revisão de 10 casos. *Rev Paul Med.* 1960; 56:352-66.
61. Vasconcelos WMP, Cardoso VM. Fígado e blastomicose sul-americana. *J Bras Med.* 1973; 1:83-90.
62. Dantas JC, Sipahi LFM. Blastomicose hepática: diagnóstico sonográfico e drenagem percutânea. *Rev Imagem.* 1985; 7(1):21-4.
63. Chaib E, de Oliveira CM, Prado PS, Santana LL, Toloi Júnior N, de Mello JB. Obstructive jaundice caused by blastomycosis of the lymph nodes around the common bile duct. *Arq Gastroenterol São Paulo.* 1988; 25(4):198-202.
64. Carmo JR, Rodrigues JHL, Adad SJ, Castilho EAS, Fischer-Puhler P. Paracoccidioidomicose: formas abdominais abordadas cirurgicamente. *Rev Med Minas Gerais.* 1991; 1(1):3-5.
65. Cesar HC, Carini A, Lauantj F, Lia N. Abdome agudo de etiologia blastomicótica. *O Hospital.* 1962; 61(3):625-37.
66. Chojniak R, Vieira RAC, Lopes A, Silva JCA, Godoy CE. Intestinal paracoccidioidomicose simulating colon cancer. *Rev Soc Bras Med Trop.* 2000; 33(3): 309-12.
67. Carvalho AS, Cerri GG, Shiroma M, Shikanai-Yasuda MA, Barone AA, Amato Neto V. Síndrome de compressão da veia cava inferior na paracoccidioidomicose. *Rev Inst Med Trop São Paulo.* 1986; 28(1):61-4.
68. Forantini OP. Blastomicose da região pancreática. *Rev Paul Med.* 1976; 31:165-72.
69. Del Negro G. Lesões das suprarrenais. In: Del Negro G, Lacaz CS, Fiorillo AM. *Paracoccidioidomicose: blastomicose sul-americana.* São Paulo: Sarvier; 1982. p. 195-202.
70. Diniz LM. Contribuição ao estudo da associação: paracoccidioidomicose: lesões suprarrenais em nosso meio. [Dissertação]. Belo Horizonte: Universidade Federal de Minas Gerais; 1985.
71. Oñate JM, Tobón AM, Restrepo A. Insuficiência suprarrenal secundária à paracoccidioidomycosis. *Biomédica.* 2002; 22:280-6.
72. Leal AMO, Magalhães PK, Martinez R, Moreira AC. Adrenocortical hormones and interleukin patterns in paracoccidioidomycosis. *J Infect Dis.* 2003; 187(1):124-7.
73. Leal AMO, Bellucci AD, Muglia VF, Lucchesi FR. Unique adrenal gland imaging features in Addison's disease caused by paracoccidioidomycosis. *AJR Am J Roentgenol.* 2003 Nov; 181(5):1433-4.
74. Oñate JM, Tobón AM, Restrepo A. Insuficiencia suprarrenal secundária à paracoccidioidomycosis. *Biomédica.* 2002; 22:280-6.
75. Marchiori E, Tannus J, Ramos RC, Cunha MLS, Pantaleão CA, Silva SC, et al. Paracoccidioidomycose das suprarrenais: avaliação por métodos radiológicos: relato de caso. *Radiol Bras.* 1990; 23:197-200.
76. Lopes DL, Araújo Sde A, Santos JP, Lyon AC, Dantas DV, Reis BS, et al. Prostatic paracoccidioidomycosis: differential diagnosis of prostate cancer. *Mem Inst Oswaldo Cruz.* 2009; 104(1):33-6.
77. Da Silva CEA, Cordeiro AF, Gollner AM. Paracoccidioidomycose do sistema nervoso central. *Arq Neuropsiq.* 2000; 58:741-7.
78. Pedroso VSP, Vilela MC, Pedroso ERP, Teixeira AL. Paracoccidioidomycosis with central nervous system involvement: review of the literature. *Rev Bras Neurol.* 2008; 44:33-40.
79. Pedroso VSP, Vilela MC, Pedroso ERP, Teixeira AL. Paracoccidioidomycosis compromising the central nervous system: a systematic review of the literature. *Rev Soc Bras Med Trop.* 2009; 42:691-7.
80. Pedroso VSP, Vilela MC, Santos PC, Cisalpino PS, Arantes RME, Rachid MA, et al. Development of a murine model of neuroparacoccidioidomycosis. *J Neuroparasitol.* 2010; 1(1):1-6.
81. Pereira WJF. Paracoccidioidomycose do sistema nervoso central – análise de 13 casos com pesquisa do antígeno gp43 por imunofluorescência. [Dissertação]. Belo Horizonte: Santa Casa de Misericórdia de Belo Horizonte; 2001.